

**Causes and mechanisms of synchronous succession trajectories in
primeval Central European mixed *Fagus sylvatica* forests**

Fabian Rey^{1,2,3}, Erika Gobet^{1,2}, Christoph Schwörer^{1,2}, Othmar Wey^{2,4}, Albert Hafner^{2,4}, Willy
Tinner^{1,2}

¹ Institute of Plant Sciences, University of Bern, Bern, Switzerland

² Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

³ Department of Environmental Sciences, University of Basel, Basel, Switzerland

⁴ Institute of Archaeological Sciences, University of Bern, Bern, Switzerland

Correspondence author

Fabian Rey, Email: fabian.rey@ips.unibe.ch

Running headline

Synchronous succession in *Fagus sylvatica* forests

Abstract

1. Natural succession trajectories of Central European forest ecosystems are poorly understood due to the absence of long-term observations and the pervasive effects of past human impacts on today's vegetation communities. This knowledge gap is significant given that currently forest ecosystems are expanding in Europe as a consequence of global change.

2. Annually laminated sediments were extracted from two small lowland lakes (Moossee 521 m a.s.l.; Burgäschisee 465 m a.s.l.) on the Swiss Plateau. We combine high-resolution palaeoecological and quantitative analyses to assess changes in vegetation during the Neolithic. We test for regionally synchronous land-use phases and plant successional patterns that may originate from complex interactions between human and climatic impacts.

3. Mixed *Fagus sylvatica* forests dominated the Swiss Plateau vegetation over millennia. During the period 6,500–4,200 cal year BP, pronounced forest disruptions accompanied by increased fire and agricultural activities occurred at c. 6,400–6,000 cal year BP, 5,750–5,550 cal year BP, around 5,400 cal year BP and at 5,100–4,600 cal year BP. Biodiversity increased during these land-use phases, likely in response to the creation of new open habitats. After decades to centuries of land-use, arboreal vegetation re-expanded. In a first succession stage, heliophilous *Corylus avellana* shrubs were replaced by pioneer *Betula* trees. These open arboreal communities were outcompeted within 150–200 years by late successional *F. sylvatica* and *Abies alba* forests. Most strikingly, cross-correlations show that these successions occurred synchronously (± 11 years) and repeatedly over large areas ($>1,000$ km²) and millennia.

4. *Synthesis*. First notable human impact shaped the primeval mixed *F. sylvatica* forests in Central Europe from c. 6,800–6,500 cal year BP on. Agrarian societies were susceptible to climate changes and we hypothesize that climate-induced, simultaneous agricultural

expansion and contraction phases resulted in synchronous regional forest successions. Currently forests are expanding in Central Europe as a result of land abandonment in marginal areas. Our results imply that mixed *F. sylvatica* forests with *A. alba* and *Quercus* may re-expand rapidly in these areas, if climate conditions will remain within the range of the mid-Holocene climatic variability (with summers c. +1–2° C warmer than today).

KEYWORDS

Biodiversity, Charcoal, Climate change, Cross-correlations, Fire ecology, Forest resilience, Non-pollen palynomorphs, Palaeoecology and land-use history, Pollen, Swiss Plateau

1. INTRODUCTION

European forest ecosystems have expanded over the last decades in response to global change involving increasing land abandonment in marginal regions (FOREST EUROPE, 2015). Most forests are still intensively managed and natural succession is restricted to a few remnant areas (Ellenberg, 2009; Kaufmann, Hauck, & Leuschner, 2018). The direction of ongoing forest change is difficult to assess since the projections are mostly based on short-term observations alone (Willis & Birks, 2006). After disturbance, forest vegetation spontaneously tends towards a late successional stage, which is often unknown because modern forests have been strongly altered by humans over centuries to millennia (Ellenberg, 2009; Puhe & Ulrich, 2001). Studying long-term successional patterns in the past (>200 years) provides unique information about vegetation dynamics under quasi-natural conditions, that may help assess future vegetation dynamics (Foster, Schoonmaker, & Pickett, 1990). Specifically, the assessment of vegetation shifts at the onset of the Neolithic, when humans gradually became sedentary in Europe (Guilaine, 2015; Müller, 2015; Schier, 2015). This period

provides insights into successional patterns of moderately disturbed forests that were only marginally shaped by human activities under environmental conditions comparable to today (Kalis, Merkt, & Wunderlich, 2003). That information is crucial for nature conservation and forest management in protected areas such as national parks (Feurdean & Willis, 2008; Lindbladh, Niklasson, Karlsson, Björkman, & Churski, 2008; Valsecchi, Carraro, Conedera, & Tinner, 2010).

Previous palaeoecological studies in Central Europe have nicely illustrated that early successional (e.g. pioneer herbs, *Corylus avellana*, *Betula*) and late successional stages (*Fagus sylvatica*, *Abies alba*) oscillated more frequently and pronouncedly after 6,500 cal year BP (Ammann, 1989; Becker et al., 2006; Kleinmann, Merkt, & Müller, 2015; Rey et al., 2017). Statistical analyses confirmed that prominent changes in vegetation were directly linked to land-use (Gobet, Tinner, Hochuli, van Leeuwen, & Ammann, 2003; Rey et al., 2013; Schwörer, Colombaroli, Kaltenrieder, Rey, & Tinner, 2015; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999). Fires were commonly used to open landscapes to create space for settlements, pastoral and arable land (Jacomet et al., 2016; Tinner, Conedera, Ammann, & Lotter, 2005), primarily disadvantaging late-successional trees and favouring the expansion of heliophilous herbs and shrubs (Conedera, Colombaroli, Tinner, Krebs, & Whitlock, 2017; Tinner et al., 2000). Many studies numerically show that land-use substantially increased open land biodiversity over the long-term (Colombaroli, Beckmann, van der Knaap, Curdy, & Tinner, 2013; Colombaroli & Tinner, 2013; Giesecke, Ammann, & Brande, 2014). However, one crucial question still remains: were the marked vegetation changes spatially synchronous? This question is important to assess the underlying causes of ecosystem change. Synchronism would speak for a superimposed forcing such as climate (i.e. precipitation and/or temperature changes) or, perhaps less likely, concerted cultural activities (Berglund, 2003; Tinner et al.,

2003), whereas diachronism would rather point to spatio-temporally randomized land-use phases of local autarchic societies that were acting independently from each other (Rösch & Lechterbeck, 2016). Until now, it was impossible to test these competing hypotheses because available records lacked temporal precision and resolution (Berglund, 2003; Tinner et al., 2003). In this study, we aim to resolve this long-standing question with high-resolution palaeoecological and time-series analyses that rely on varved sediment chronologies reaching exceptional temporal precision of c. ± 20 –50 cal years (Rey et al., 2018). The overarching goal of the study is to explore the nature of re-occurring broad-scale successional patterns in mixed beech forests after moderate human impact to refine existing projections of future forest transformations (Bugmann et al., 2015; Ruosch et al., 2016) under global change conditions that are currently triggering forest expansions in Central Europe and adjacent regions.

2. MATERIAL AND METHODS

2.1. Study sites

The study area is located on the western Swiss Plateau in southern Central Europe (Figure 1). Moossee is a small lake close to the city of Bern at 521 m a.s.l. (47°1'17.0"N, 7°29'1.7"E). The lake has a maximum water depth of 22 meters and a surface area of 0.31 km². Burgäschisee is a small lake further to the northeast (c. 25 km from Moossee) at 465 m a.s.l. (47°10'8.5"N, 7°40'5.9"E). It has a maximum water depth of 31 meters and a surface area of 0.21 km². Both lakes are eutrophic today with anoxic waters in the hypolimnion (Guthruf, Zeh, & Guthruf-Seiler, 1999). The present climate is temperate with mean annual temperatures of 8.8°C and an annual rainfall of 1,059 mm at Moossee (data from Bern/Zollikofen at c. 3 km distance, MeteoSwiss, 2017). At Burgäschisee, it is slightly warmer and wetter with mean annual temperatures of 9.1°C and an annual rainfall of 1,088 mm (data from Koppigen, at c. 6 km

distance, MeteoSwiss, 2017). The current vegetation at both lakes is dominated by mixed *F. sylvatica* forests on the surrounding hills and scattered mixed *Alnus glutinosa*-*Fraxinus excelsior* stands on the lake shores. Most of the flat areas around the lakes are intensively used for agriculture. The region has been important for agricultural activities for millennia as indicated by frequent local archaeological findings dating back to at least 6,800-6,500 cal year BP (Hafner, Harb, Amstutz, Francuz, & Moll-Dau, 2012; Harb, 2017; Wey, 2012).

2.2. Corings and chronologies

In 2014, three parallel sediment cores at Moossee and two parallel sediment cores at Burgäschisee were retrieved with a UWITEC piston corer (core diameter: 9 cm, core length: 200 cm) in deep parts of the lakes. A total of 7 m of lake sediments were sampled at each site covering c. 9,000 years. For this study, we focus on the Neolithic to Early Bronze Age period (6,500–3,800 cal year BP) during which the sediments are annually laminated (i.e. varved) at both lakes (see Rey et al., 2018).

The Moossee chronology is based on 27 terrestrial plant macrofossils and varve counts (Rey et al., 2018). The program OxCal 4.3 (V-sequence; Bronk Ramsey, 1994, 1995, 2001; Bronk Ramsey, van der Plicht, & Weninger, 2001) and the IntCal13 calibration curve (Reimer et al., 2013) were used to estimate the age-depth model (Figure 2a). The Burgäschisee chronology is based on 22 terrestrial plant macrofossils and basic assumptions about the sedimentation rates from a previous chronology (Rey et al., 2017, 2018). Here, the program OxCal 4.3 (U/P-sequences; Bronk Ramsey, 1994, 1995, 2001, 2008; Bronk Ramsey & Lee, 2013) and the IntCal13 calibration curve (Reimer et al., 2013) were used for the estimations of the age-depth model (Figure 2b). Both records indicate fairly regular sedimentation rates for the investigated time interval with an average of 26 years/cm at Moossee and 18.7 years/cm at Burgäschisee.

138 **2.3. Pollen, macrofossil, non-pollen palynomorphs and charcoal analyses**

139 All palaeoecological analyses were conducted on the same cores and for the same sample
140 depths (Moossee: 312 samples, Burgäschisee: 308 samples). We applied a contiguous,
141 approximately constant year sampling (10 ± 2 years/sample). For all microscopic analyses
142 (pollen, non-pollen palynomorphs (NPPs), microscopic charcoal), samples of 1 cm^3 were
143 treated with HCl, KOH, HF, acetolysis, sieved with a mesh size of $500 \mu\text{m}$ and mounted in
144 glycerine following standard palynological methods (Moore, Webb, & Collinson, 1991).
145 *Lycopodium* tablets (University of Lund batch no. 1031 with $20,848 \pm 3,457$ spores per tablet
146 for Moossee and University of Lund batch no. 177745 with $18,584 \pm 1,853$ spores per tablet
147 for Burgäschisee) were added prior to the chemical treatment to estimate microfossil
148 concentrations (Stockmarr, 1971). We used palynological keys (Beug, 2004; Moore, Webb, &
149 Collinson, 1991), photo atlases (Reille, 1992) and the reference collection at the Institute of
150 Plant Sciences (University of Bern) to identify pollen and spores under a light microscope at
151 $400\times$ magnification. Phase contrast under $1,000\times$ magnification was used to separate various
152 Cerealia-types (*Hordeum*-type, *Triticum*-type; Beug, 2004). The minimum number of
153 terrestrial pollen grains counted per sample was 500 with a total of 132 (Moossee) and 113
154 (Burgäschisee) identified terrestrial pollen types. The results are presented as percentages of
155 the terrestrial pollen sum (Figures 3 and 4), in which we excluded spores, pollen of aquatic
156 plants and other microfossils (e.g. stomata, algae cells). Pollen can be dispersed by wind over
157 large distances (up to 50 km) and generally represents extra-local to regional vegetation
158 dynamics (Conedera, Tinner, Cramer, Torriani, & Herold, 2006). However, since the sites are
159 rather small (Moossee: 0.31 km^2 , Burgäschisee 0.21 km^2), we assume that the pollen source

area only spans several hundred meters to at most a few kilometers (Conedera, Tinner, Cramer, Torriani, & Herold, 2006; Sugita, 1994).

Samples for macrofossil analyses (including macroscopic charcoal) had mean volumes of 17.5 cm³ (Moossee) and 11 cm³ (Burgäschisee) and were sieved with a mesh size of 200 µm. Terrestrial remains (e.g. *F. sylvatica* bud scales) were identified under a stereo microscope. Macrofossil concentration (remains cm⁻³) were calculated and plotted in Figures 3 and 4. Plant macrofossils mainly indicate local vegetation as a result of short dispersal distances (1–100 m; Birks, 2003).

Microscopic charcoal particles >10 µm and <500 µm were counted on pollen slides following Tinner and Hu (2003) and Finsinger and Tinner (2005) and presented as microscopic charcoal influx values (particles cm⁻² year⁻¹; Figures 3 and 4). These values were used as a proxy for regional fire activity since microscopic charcoal particles are dispersed over distances of c. 20–50 km (Adolf et al. 2018; Tinner et al. 1998).

Macroscopic charcoal pieces >600 µm were identified under a stereo microscope during macrofossil analysis. Their occurrence (pieces cm⁻³) may primarily reflect local fire activity within few hundred meters distance (Adolf et al., 2018; Figures 3 and 4). This procedure (large sampling volume, size threshold >600 µm) allows reducing the influence of regional fires on macroscopic charcoal, without using de-trending and peak isolation approaches (Adolf et al., 2018).

Green algae (*Botryococcus*, *Tetraedron*, *Coelastrum*) and akinetes of cyanobacteria (*Anabaena*, *Aphanizomenon*) were counted on pollen slides to estimate changes in the trophic levels. Although, single NPP taxa might occur under natural conditions, contemporaneous or consecutive high numbers of green algae and cyanobacteria are considered as a proxy for eutrophication during or after local anthropogenic impacts (van Geel, Mur, Ralska-

Jasiewiczowa, & Goslar, 1994; Hillbrand, van Geel, Hasenfratz, Hadorn, & Haas, 2014). All NPPs are presented as percentages of the terrestrial pollen sum (Figures 3 and 4).

Local pollen assemblage zones (LPAZ; Figures 3 and 4) were delimited using optimal sum-of-squares partitioning (Birks & Gordon, 1985) with the program ZONE 1.2 (Juggins, 1991). Statistically significant zones were assessed with the program BSTICK (Line & Birks, unpublished software) following the broken-stick method (Bennett, 1996). The programs Tilia 2.0.41 and CorelDraw were used to plot the data for both high-resolution sequences (Figures 3 and 4).

2.4. Biodiversity estimations

Rarefaction analysis was used to calculate the palynological richness (PRI), which is commonly used as a proxy for species richness at local to regional scale in many palaeoecological studies (e.g. Birks & Line, 1992; Colombaroli & Tinner, 2013; Odgaard, 1999; Schwörer, Colombaroli, Kaltenrieder, Rey, & Tinner, 2015). Rarefaction analysis allows estimating the number of taxa per sample for a constant counting sum (Birks & Line, 1992), which was 500 at both sites. The probability of interspecific encounter (PIE; Hurlbert, 1971) was used as a measure of palynological evenness. In previous studies PIE was used to evaluate effects of few dominant pollen producers (e.g. *C. avellana*; van der Knaap, 2009) on the palynological assemblages and in particular on palynological richness. To assess such effects, PIE-derived evenness-detrended palynological richness (DE-PRI) was calculated following Colombaroli, Beckmann, van der Knaap, Curdy, and Tinner (2013). This approach aims at removing the evenness trend from palynological richness by building an ordinary least square regression (OLS) between the dependent (PRI) and the independent variable (palynological evenness). The resulting residuals of pollen richness (= PRI - PIE) are then distributed around the original pollen richness

values (Colombaroli & Tinner, 2013). Only if PRI and DE-PRI show similar changes and trends, we assume that the species richness estimates are unaffected by evenness effects. The DE-PRI approach has proven to be valuable to estimate contemporary species richness patterns at the European scale (Adolf, 2017). The program R statistics (R Development Core Team, 2016) was used for all calculations (see results in Figures 3 and 4).

2.5. Time-series analyses (cross-correlations)

Cross-correlations (Gobet, Tinner, Hochuli, van Leeuwen, & Ammann, 2003; Green, 1981; Rey et al., 2013; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999) were calculated to identify leads and lags between fire (microscopic charcoal influx values) and vegetation (pollen percentages, PRI, PIE, DE-PRI) by using the program MYSTAT 12 (Systat, 2007). The time window for all calculations was 6,500–4,150 cal year BP that included 209 samples at Moossee (1 sample = 11.1 ± 0.2 cal years) and 266 samples at Burgäschisee (1 sample = 8.7 ± 0.6 cal years). Cross-correlation coefficients for microscopic charcoal influx values vs. pollen percentages, PRI, PIE and DE-PRI (see Figure 5) were calculated at ± 50 lags corresponding to ± 555 cal years (Moossee) and ± 435 cal years (Burgäschisee), following Bahrenberg, Giese, Mevenkamp, and Nipper (2008), with maximum lag numbers corresponding to one fourth of the sample number N (lag number $\leq N/4$). The correlation coefficients at lag 0 between microscopic charcoal and selected pollen types are represented as detailed correlograms (Figure 6). For cross-correlation analyses between the two sites (e.g. *F. sylvatica* percentages at Moossee vs. *F. sylvatica* percentages at Burgäschisee), the time series with higher time resolution (Burgäschisee) was adjusted to the one with lower resolution (Moossee) by combining neighbouring samples following Tinner et al. (2015). This led to a total of 209 samples (1 sample = 1 lag = 11.1 ± 0.2 cal years) and ± 50 lags (= ± 555 cal years) that were

included in the cross-correlation calculations (Figure 7). Variables were not de-trended prior to the cross-correlation analyses because the vegetation type remains more or less uniform for the whole time window without pronounced population trends (see Figures 3 and 4; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999). The 95% (2σ) confidence intervals of the correlations were estimated by computing ± 2 standard errors of the correlation coefficients (Bahrenberg, Giese, Mevenkamp, & Nipper, 2008). This corresponds to a two-sided significance level (α) of 5% (Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999).

3. RESULTS

3.1. Vegetational, agricultural and fire dynamics

Both studied pollen records (Moossee: 7,000–3,900 cal year BP, Burgäschisee: 6,500–3,800 cal year BP) have a high temporal resolution of c. 10 cal years per sample and an extraordinary chronological precision (Rey et al., 2018). Zonation analyses show a high number of significant LPAZ (Moossee: 15, Burgäschisee: 14; see Figures 3 and 4), suggesting high variability of vegetation composition over time. In general, *F. sylvatica* was dominant in the closed forests (pollen percentages generally $\geq 20\%$). This pollen-based interpretation is supported by regular finds of macrofossils (bud scales). Other important taxa in the forests included *Quercus* and *A. alba*. The latter shows slightly higher pollen percentages at Burgäschisee (up to 10% instead of 5%), probably as a result of increasing precipitation towards the east of the Swiss Plateau, making the species more competitive under mesophilous conditions (Rey et al., 2017). However, macrofossil and stomata finds of *A. alba* point to local occurrences of the species at both sites. *Alnus glutinosa* was preferably growing on the wet soils at the lake shores as indicated by high and fairly stable pollen percentages (around 25%) and abundant macrofossils finds (e.g. *A. glutinosa* fruits). Pollen percentages of other subordinate taxa such

256 as *Tilia*, *Ulmus* and *Taxus baccata* steadily decrease until c. 5,000 cal year BP after which they
257 remain low (<2%, see *Ulmus*) or even disappear (see *Tilia* and *T. baccata*), most likely as a
258 result of overexploitation and fire disturbance (Rey et al., 2017; Tinner, Conedera, Ammann,
259 & Lotter, 2005). Distinct phases with low *F. sylvatica* pollen percentages (<10%), increased
260 cultural indicators (Cerealia-type, *Plantago lanceolata*, *Linum usitatissimum*-type), apophytes
261 (*Urtica*), other herbs (Poaceae, *Allium*-type) and heliophilous shrubs (*C. avellana*, *Juniperus*)
262 are apparent at both sites. A first phase is dated at 6,550–6,150 cal year BP at Moossee and
263 at 6,300–5,950 cal year BP at Burgäschisee. Subsequently, similar phases occurred at 5,750–
264 5,550 cal year BP, at around 5,400 cal year BP and at 5,100–4,600 cal year BP at both sites.
265 Finally, a last phase is dated around 3,900 cal year BP at Moossee. Periods of forest disruption
266 were generally accompanied by high regional fire activity as suggested by increasing
267 microscopic charcoal influx values (>5,000 particles cm⁻² yr⁻¹), sometimes overlapping with
268 local fire activity as inferred from macroscopic charcoal presence (≥600 µm). This finding is in
269 good agreement with local archaeological evidence (e.g. pile dwellings, light grey bars in
270 Figures 3 and 4). Some settlements around the lake had impacts on the water quality as shown
271 by high numbers of cyanobacteria (*Anabaena*, *Aphanizomenon*) and green algae (*Tetraedron*)
272 between 5,750 and 5,550 cal year BP at both lakes and around 4,600 cal year BP at Moossee.
273 Such blooms are best explained by strong lake water eutrophication. After the settlement
274 phases, *Betula* as a pioneer tree species quickly recolonized abandoned places (pollen
275 percentages up to 30% and abundant macrofossils). After intermediate phases often with *F.*
276 *excelsior*, *Ulmus* and *Tilia* expansions, these trees were unremittingly replaced within 150–200
277 years by late successional tree species (*F. sylvatica*, *A. alba*) if no further disturbances
278 occurred. These succession cycles were simultaneously repeated at both sites and are not only

visible in the pollen percentage data but are also confirmed by plant macrofossils, indicating possible broad-scale succession trajectories.

3.2. Biodiversity reconstruction

Palynological richness (PRI) and evenness-detrended palynological richness (DE-PRI) show good agreements (Figures 3 and 4), suggesting that overall trends in palynological richness are not strongly affected by evenness. PRI and DE-PRI generally increase during phases with higher human impact (>25 pollen types per sample), i.e. around 6,400 cal year BP, 4,600 cal year BP and 3,900 cal year BP at Moossee, at 5,700–5,600 cal year BP at both sites, and around 4,900 cal year BP at Burgäschisee. These increases are directly related to vegetation openings and the introduction of cultivated plants (*Hordeum*-type, *Triticum*-type, *L. usitastissimum*-type). The establishment of open land for Neolithic farming created new habitats for short-lived weeds (*P. lanceolata*), apophytes (*Urtica*) and light-demanding shrubs (*C. avellana*, *Sambucus nigra*). Palynological evenness as inferred from PIE (Figures 3 and 4) is more or less stable (PIE around 0.85). Slightly lower values of palynological evenness (PIE around 0.8) are recorded in phases when either *Betula* or *F. sylvatica* pollen grains are dominant and the forests were rather closed, suggesting that the expansion of monospecific stands may have affected species evenness. On the other hand, phases with open forests (e.g. 5,700–5,600 cal year BP at both sites) have usually high palynological evenness values (PIE ≥ 0.85), suggesting that vegetation evenness is correlated to openness.

3.3. Time-series analyses

At both sites, pollen percentages of trees have significant negative correlations with microscopic charcoal influx (maximum negative correlations at lag 0; Figure 5). *F. excelsior* is

303 a good example to illustrate immediate negative fire impacts. The negative correlations of *F.*
304 *sylvatica* pollen percentages is, however, markedly delayed at both sites (c. 320 years after a
305 fire at Moossee respectively c. 80 years after a fire at Burgäschisee), possibly indicating a
306 slightly higher fire tolerance compared to other tree species (Tinner et al., 2000). On the other
307 hand, pollen percentages of herbs (e.g. *Cerealia*-type) reach maximum positive correlations
308 with microscopic charcoal at lag 0, showing that agropastoral activities were strongly
309 connected to the use of fire. Similarly, correlations between microscopic charcoal influx and
310 PRI, DE-PRI and at Moossee also PIE suggest that biodiversity increased in the new habitats
311 (i.e. open areas) created by anthropogenic burning for agricultural purposes. Pollen
312 percentages of shrubs such as *C. avellana* and *Salix* also have significant positive correlations,
313 e.g. at Burgäschisee maximum positive correlations are reached at lag +4 for *C. avellana* (c. 35
314 years after a fire) and at lag 0 for *Salix* (Figure 5b). *Betula* shows maximum positive correlations
315 at lag +35 (c. 390 years after a fire; Figure 5a) at Burgäschisee and at lag +41 (c. 355 years after
316 a fire; Figure 5b) at Moossee. This result with *C. avellana* preceding *Betula* pollen percentages
317 is likely caused by the successional trajectories after fire disturbance. Later on, the recovery
318 of *F. excelsior* follows (lag +41 and lag +45, corresponding to c. 455 and c. 390 years) after the
319 fire-related decline (negative correlations) as well as *F. sylvatica* (lag +50 = c. 435 years; Figure
320 5b). Correlation coefficients at lag 0 between microscopic charcoal influx values and pollen
321 types or PRI, PIE and DE-PRI show similar linkages (Figure 6), with significant positive
322 correlations for cultural indicators (*Hordeum*-type, *Triticum*-type), other herbs (e.g. *Allium*-
323 type, *Mentha*-type, *Papaver rhoeas*-type, *Urtica*), PRI, DE-PRI and shrubs (e.g. *C. avellana*,
324 *Salix*, *S. nigra*). In contrast, most tree pollen types (e.g. *Quercus*, *F. excelsior*, *F. sylvatica*,
325 *Ulmus*, *T. baccata*) have significant negative correlations with microscopic charcoal,
326 suggesting that forest trees were most affected by anthropogenic burning. However, A.

glutinosa-type as an important tree pollen type is indifferent showing neither significant positive nor negative correlations with charcoal-inferred fire incidence.

If cross-correlations are used to compare the two sites, a striking pattern appears (Figure 7). Total shrubs and trees as well as some taxa (e.g. *C. avellana*, *Betula*, *F. sylvatica*, *Tilia*) show pronounced positive correlations with maximum correlations at lag 0, suggesting synchronous oscillations within the pooled sample resolution (11.1 ± 0.2 cal years). Cross-correlations between pollen percentages of herbs (Moossee data vs. Burgäschisee data), *F. excelsior* and *A. alba* only slightly differ and may reflect local vegetation variability, however, significant positive correlations also center around lag 0 with maximum positive correlations within c. ± 35 years. Of particular interest are the correlations for the cereals, that also show this pattern, confirming that the cultural phases at the two sites were largely coeval with only minor differences, possibly indicating positive synergy effects of cereal production in old and new prehistoric centres. Similarly, also PRI and DE-PRI co-vary in time at the two sites. Taken together, these results quantitatively show that the general succession patterns after disturbance were not only similar as presented in Figures 3 and 4 but in fact synchronous at the two sites.

4. DISCUSSION

4.1. Impacts of land-use on temperate forests and synchronous patterns of forest succession after disturbances

The palaeobotanical data suggest the dominance of mixed beech forests over millennia at our two study sites Burgäschisee and Moossee. Land-use phases contributed to openings and a gradual conversion from mixed beech to almost pure beech forests. These general patterns have been observed elsewhere in the lowlands of southern Central Europe after c. 7,000 cal

year BP (e.g. Ammann, 1989; Kleinmann, Merkt, & Müller, 2015; Rösch & Lechterbeck, 2015; Tinner & Ammann, 2005; Tinner, Conedera, Ammann, & Lotter, 2005). However, local differences in species composition resulted from dissimilar environmental conditions. For instance in Switzerland, moisture availability increases towards higher elevations and to the east. This moisture gradient may have led to the co-dominance of *A. alba* in the forests (Gobet & Tinner, 2012; van der Knaap, van Leeuwen, & Ammann, 2004; Lotter, 1999; Wehrli, Tinner, & Ammann, 2007), given that as the most shade tolerant and tallest tree species of Europe, silver fir is particularly competitive under mesophilous conditions (Ellenberg, 2009; Ruosch et al., 2016; Tinner et al., 2013). Conversely, drier conditions in the Lake District (Seeland) in the rain shadow of the Jura Mountains may have promoted *Quercus* spp. including submediterranean *Q. pubescens* (Ammann, 1989; Hadorn, 1992; Steiger, 2010).

Several diebacks of mixed *F. sylvatica* forests occurred between 6,400 and 6,000 cal year BP, at 5,800–5,650 cal year BP, 5,400–5,300 cal year BP, at 5,100–4,700 cal year BP at both sites and around 3,900 cal year BP at Moossee only. The combined charcoal and pollen evidence as well as the cross-correlation analyses suggest that burning was used as a tool for opening the mixed *F. sylvatica* forests, creating a mosaic of new habitats that advantaged many herbaceous species, increasing the overall biodiversity (see biodiversity estimations in Figures 3 and 4). A similar pattern has been reconstructed in previous studies and might be typical for European landscapes (Colombaroli, Beckmann, van der Knaap, Curdy, & Tinner, 2013; Colombaroli & Tinner, 2013; Giesecke, Ammann, & Brande, 2014). Prehistoric slash-and-burn activities were strong enough to release a wide-scale expansion of early-successional shrublands, typically dominated by *C. avellana*, *Betula* and *Alnus* (Aaby, 1986; Tinner, Conedera, Ammann, & Lotter, 2005; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999). In contrast, most tree species were strongly reduced by fires and other human disturbances

such as logging, pollarding and browsing (e.g. Haas, Karg, & Rasmussen, 1998; Haas & Schweingruber, 1993; Pott, 1985; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999). Most interestingly, some tree species such as *F. sylvatica* and *A. alba* seem to have fully recovered and even expanded after forest disruptions (within c. 150–200 years after a land-use phase), suggesting long-term resilience to moderate human disturbance, while others (*Ulmus*, *Tilia*, *Acer* and *T. baccata*) markedly diminished, suggesting high sensitivity of these species. Ultimately, at c. 5,000 cal year BP (Figure 3 and 4) this process of repeated disturbances led to a strong impoverishment in species diversity of forest ecosystems (Tinner & Ammann, 2005), generating the advent of monospecific forests that are characteristic for Central European landscapes today (e.g. almost pure spruce, beech, larch, oak stands, depending on altitude and other environmental conditions; Ellenberg, 2009; Gobet, Vescovi, & Tinner, 2010).

Here, we quantitatively demonstrate for the first time that forest succession cycles over 3,000 years were regionally synchronous (Figure 7). The distance between the two sites (c. 25 km) exceeds the expected pollen catchment expected for the two study sites (most pollen from within a few hundred meters to at most a few kilometers; Conedera, Tinner, Cramer, Torriani, & Herold, 2006), arguing for coeval population dynamics over the millennia within the time resolution and precision available (c. 20–50 cal years; Rey et al., 2018). Because this pattern was found in the cross-correlation analyses for several tree taxa as well as for the tree percentage sum, we consider it to be very reliable. Synchronous forest disruptions (tree pollen <80%) may have occurred at several distant sites in Central and Southern Europe (Figure 8). The few available high-resolution (c. 10–20 years) and high-precision (sufficient radiocarbon dates on terrestrial material) time series covering the entire Neolithic and the onset of the Bronze Age reveal coeval tree population reductions at c. 6,400–

5,900 cal year BP, 5,750–5,600 cal year BP, 5,400 cal year BP, 5,100–4,700 cal year BP and around 3,900 cal year BP (Ammann, 1989; Kleinmann, Merkt, & Müller, 2015; Lotter, 1999; Rösch, Kleinmann, Lechterbeck, & Wick, 2014a, b; Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999; Wehrli, Tinner, & Ammann, 2007). One of these sites, Lago di Origlio, is located in the lowlands south of the Alps (Sottoceneri), where environmental conditions are typical for submediterranean Southern Europe (warmer summers, milder winters). We thus hypothesize the existence of supra-regional synchronicity of forest disturbances during the Neolithic, as it was formerly advocated for Central and Southern Europe (including Sicily) for the Bronze Age, the Iron Age, the Roman Period and the subsequent early medieval times (Finsinger & Tinner, 2006; Gobet, Tinner, Hochuli, van Leeuwen, & Ammann, 2003; Tinner et al., 2003, 2009).

4.2. Possible impacts of climate on land-use phases

Simultaneous broad-scale deforestation phases over large areas caused by land-use may mainly reflect superimposed causes such as climate change (e.g. Berglund, 2003; Tinner et al., 2003, 2009). Our study sites showing coeval vegetation succession patterns are both located in landscapes providing fertile arable lands (Guthruf, Zeh, & Guthruf-Seiler, 1999). They are mostly lacking major in- and outflows, thus settlements around the lakes, especially at the shores (e.g. pile dwellings) may have been affected by lake level changes (Guthruf, Zeh, & Guthruf-Seiler, 1999). We here assume that people settled at lower elevations preferably near lake shores/rivers during warm and dry phases and subsequently moved upwards to slightly higher elevations during cool and wet phases. Thus, the more densely populated warm and dry phases might be preferentially reflected in the archaeological and dendrochronological records e.g. at Lake Biel (Fischer, Hafner, Stapfer, Marti, & Affolter, 2017; Hafner & Suter,

2000; Stöckli, 2016; Suter, 2017) and Lake Zurich (Bleicher et al., 2017), whereas the fewer settlements on dry soils during cool and wet phases (e.g. hilltop sites) were more exposed to erosion, hence artifacts are scarce.

Common land-use phases across Central and Southern Europe were possibly driven by temperature oscillations, which were usually associated with moisture changes. After 8,200 cal year BP, warm periods in and around the Alps were generally associated with dry conditions, while decadal to centennial scale cold excursions were wetter, comparable to the conditions during the Little Ice Age (Haas, Richoz, Tinner, & Wick, 1998; Tinner & Lotter, 2006). Indeed wet phases, as reconstructed from lake-level proxies occurred at around 6,300, 5,500 and 4,200 cal year BP (Magny, 2004, 2013), when crop production around the lakes was low. The linkage to warm phases as reconstructed from tree-ring records (Bircher, 1982, 1986; Renner, 1982) and the Greenland stable oxygen isotope records (Vinther et al., 2006) is less pronounced (Figure 9). Although the Swiss tree-ring records of temperature change are well in line with the Greenland stable oxygen isotope record, the chronological precision of these records (c. ± 50 –100 cal years) is lower than that of the dendrochronologically dated archaeological evidence (c. ± 5 –10 years). Previous studies have emphasized the good match between the dendrochronologically dated archaeological phases and solar activity, as reconstructed from the ^{14}C -residuals at multiannual to multidecadal scales (Gross-Klee & Maise, 1997; Kleinmann, Merkt, & Müller, 2015; Magny, 2004, 2013). The link between solar activity and archaeological evidence becomes again evident when comparing the land-use phases at both study sites with the reconstructed average total solar irradiance (TSI; Steinhilber, Beer, & Fröhlich, 2009; see Figure 9). Periods with high solar irradiance corresponded with rather dry conditions (low lake levels) and increases of cultural indicators (sum of cereals, *L. usitatissimum*, *P. lanceolata*), local–regional fires (macroscopic and

microscopic charcoal) and biodiversity (PRI). Conversely, periods with lower TSI and wetter conditions (high lake levels) corresponded to land abandonment phases. We assume that under adverse wet and cool conditions agricultural yields collapsed and human population densities declined over large areas, as shown by the excellent agreement across distant sites (Figure 8). Solar irradiance may have additionally advantaged crop production, given that the cereals, which were the main source of calories, originate from the Near East and are adapted to high insolation.

4.3. Implications for global change ecology

Our study shows that over the millennia temperate forests dominated by *F. sylvatica* and *A. alba* were resilient to prehistoric human impacts in the Swiss Plateau. Forest succession after disturbances generally started with light-loving pioneer species (*C. avellana*, *Betula*) and was completed within 150–200 years by late successional mixed beech forests. Such palaeoecological information might be crucial to develop sustainable strategies for nature conservation and forest management (Feurdean & Willis, 2008; Lindbladh, Niklasson, Karlsson, Björkman, & Churski, 2008; Valsecchi, Carraro, Conedera, & Tinner, 2010), specifically under global-change conditions that are currently releasing forest ecosystem expansions in abandoned rural areas (FOREST EUROPE, 2015). We assume that *F. sylvatica* forests will remain key communities in Central Europe, if future (summer-) climate warming does not exceed the Holocene variability range (c. +1–2°C compared to the 20th century; Heiri, Tinner, & Lotter, 2004). Other trees such as *F. excelsior*, *Tilia* and *Ulmus* might also re-expand under low disturbance levels (e.g. no browsing, no pollarding) provided that no further devastating diebacks due to pathogenic diseases (e.g. *Ulmus*, *F. excelsior*) occur (see Brasier, 1991; McKinney, Nielsen, Hansen, & Kjær, 2011). Anticipated climate change (Appenzeller et

al., 2011; Kovats et al., 2014), however, with generally lower annual precipitation rates may markedly disadvantage *F. sylvatica* and promote *A. alba* and *Quercus* spp. (including submediterranean *Q. pubescens*) that are less-drought sensitive and thus more competitive under drier conditions (Bugmann et al., 2015; Ellenberg, 2009; Ruosch et al., 2016; Tinner et al., 2013). Predicted higher risk of forest fires (Kovats et al., 2014) may affect fire-sensitive *A. alba* (Tinner et al., 2013), or conversely *A. alba* and the expansion of mediterranean evergreen trees (e.g. *Quercus ilex*) may further reduce fire incidence because of positive feedbacks mainly related to decreased biomass flammability (Henne et al., 2015, 2018).

5. CONCLUSIONS

We numerically demonstrate for the first time that vegetation dynamics at two distant sites were synchronous over more than 2,000 years, with a pooled sampling resolution of 11 years. Interactions between the first agrarian societies and their environment were complex, with humans significantly shaping vegetation structure (creation of open land, shrublands) and composition (shift in species). Conversely, the rise and fall of early farming societies was likely dependent on climate. Favourable climatic conditions (i.e. warm and dry summers) probably led to an increase in agricultural yields, the expansion of farming activities and resulting forest openings, whereas unfavourable climatic conditions (i.e. cold and wet summers) likely caused crop failures, abandonment of agricultural areas and forest succession. A better understanding of the environmental and societal factors controlling coeval land-use dynamics as shown in this study would require new climate proxy data (e.g. temperature reconstruction from well dated and complete Holocene tree ring series). On the basis of our results and considering the ongoing spread of temperate forests in lowland Central Europe, we conclude that the existing beech forest ecosystems are resilient to anthropogenic

disturbances under a changing climate, if the amplitude does not exceed the range of Holocene climate variability. Our results also indicate that abandoned agricultural land can revert to quasi-natural mixed beech forests within 150–200 years. Climate warming exceeding the Holocene variability range, however, has been identified as an important threshold that might cause the rapid collapse of beech forests, giving way to novel communities that are today restricted to Southern Europe.

ACKNOWLEDGEMENTS

Willi Tanner, Julia Rhiner, Sandra O. Brügger, Lena Thöle, Petra Beffa, Daniele Colombaroli, Claire Rambeau and Stéphanie Samartin helped with the corings. We thank Sönke Szidat and Edith Vogel for conducting the radiocarbon dating, André F. Lotter for sharing his knowledge about varved lake sediments, Werner E. Stöckli for fruitful discussions, Christian Harb for providing the archaeological data at Moossee and Jacqueline FN van Leeuwen for helping with pollen identifications. Special thanks go to Crystal H. McMichael, H. John B. Birks and one anonymous reviewer for their valuable comments on a previous version of the manuscript. Funding was provided by the Swiss National Science Foundation (SNF 200021_149203/1).

AUTHORS' CONTRIBUTIONS

F.R. performed all pollen, macrofossil and quantitative analyses, figure construction, and led the writing; E.G. and W.T. initiated and designed the research project and wrote substantial parts of the manuscript; C.S. helped to conduct the biodiversity estimations; A.H. and O.W. provided archaeological data; W.T. and A.H. obtained funding. All authors contributed critically to the drafts and gave final approval for publication.

519 **DATA ACCESSIBILITY**

520 The data will be available through the Alpine Palynological Database (ALPADABA) after
521 publication.

522 REFERENCES

- 523 Aaby, B. (1986). Trees as anthropogenic indicators in regional pollen diagrams from eastern Denmark.
 524 In K. E. Behre (Ed.), *Anthropogenic indicators in pollen diagrams* (pp. 73–93). Rotterdam, The
 525 Netherlands: Balkema.
- 526 Adolf, C. (2017). *Calibrating fire and vegetation proxies across large geographic and climatic gradients*
 527 *for quantitative environmental reconstructions*. Dissertation, University of Bern, Switzerland.
- 528 Adolf, C., Wunderle, C., Colombaroli, C., Weber, H., Gobet, E., Heiri, O., ... Tinner, W. (2018). The
 529 sedimentary and remote-sensing reflection of biomass burning in Europe. *Global Ecology and*
 530 *Biogeography*, 27, 199–212. <https://doi.org/10.1111/geb.12682>
- 531 Ammann, B. (1989). Late-Quaternary palynology at Lobsigensee. Regional vegetation history and local
 532 lake development. *Dissertationes Botanicae*, 137, 1–157.
- 533 Appenzeller, C., Bey, I., Croci Maspoli, M., Fuhrer, J., Knutti, R., Kull, C., & Schär, C. (2011). Swiss climate
 534 change scenarios CH2011. Zurich, Switzerland: C2SM, MeteoSwiss, ETH, NCCR Climate, OcCC.
- 535 Bahrenberg, G., Giese, E., Mevenkamp, N., & Nipper, J. (2008). *Statistische Methoden in der*
 536 *Geographie. Band 2: Multivariate Statistik*. Stuttgart, Germany: Borntraeger.
- 537 Becker, A., Ammann, B., Anselmetti, F. S., Hirt, A. M., Magny, M., Rachoud, A.-M., ... Wüthrich, C.
 538 (2006). Paleoenvironmental studies on Lake Bergsee, Black Forest, Germany. *Neues Jahrbuch für*
 539 *Geologie und Paläontologie - Abhandlungen*, 240, 405–445.
- 540 Bennett, K. D. (1996). Determination of the number of zones in a biostratigraphical sequence. *New*
 541 *Phytologist*, 132, 155–170. <https://doi.org/10.1111/j.1469-8137.1996.tb04521.x>
- 542 Berglund, B. E. (2003). Human impact and climate changes – synchronous events and a causal link?
 543 *Quaternary International*, 105, 7–12. [https://doi.org/10.1016/S1040-6182\(02\)00144-1](https://doi.org/10.1016/S1040-6182(02)00144-1)
- 544 Beug, H.-J. (2004). *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*.
 545 Munich, Germany: Pfeil.
- 546 Bircher, W. (1982). *Zur Gletscher- und Klimageschichte des Saastales. Glazialmorphologische und*
 547 *dendroklimatologische Untersuchungen*. Physische Geographie, Vol. 9. University of Zurich,
 548 Switzerland.
- 549 Bircher, W. (1986). Dendroecology applied in mountain regions. In B. E. Berglund (Ed.), *Handbook of*
 550 *Holocene palaeoecology and palaeohydrology* (pp. 387–403). Chichester, UK: Wiley.
- 551 Birks, H. H. (2003). The importance of plant macrofossils in the reconstruction of Lateglacial vegetation
 552 and climate: examples from Scotland, western Norway, and Minnesota, USA. *Quaternary Science*
 553 *Reviews*, 22, 453–473. [https://doi.org/10.1016/S0277-3791\(02\)00248-2](https://doi.org/10.1016/S0277-3791(02)00248-2)
- 554 Birks, H. J. B., & Gordon, A. D. (1985). *Numerical methods in quaternary pollen analysis*. London, UK:
 555 Academic Press.
- 556 Birks, H. J. B., & Line, J. M. (1992). The use of rarefaction analysis for estimating palynological richness
 557 from Quaternary pollen-analytical data. *The Holocene*, 2, 1–10.
 558 <https://doi.org/10.1177/095968369200200101>
- 559 Bleicher, N., Antolín, F., Gobet, E., Jacomet, S., Tinner, W., Hüster Plogmann, H., & Häberle, S. (2017).
 560 Umwelt-, Landschaft- und Besiedlungsgeschichte. In N. Bleicher N & C. Harb (Eds.), *Zürich-*

561 *Parkhaus Opéra. Eine neolithische Feuchtbodenfundstelle. Band 3: Naturwissenschaftliche*
562 *Analysen und Synthese* (pp. 226–238). Monographien der Kantonsarchäologie Zürich 50.

563 Brasier, C. M. (1991). *Ophiostoma novo-ulmi* sp. nov., causative agent of current Dutch elm disease
564 pandemics. *Mycopathologia*, 115, 151–161.

565 Bronk Ramsey, C. (1994). Analysis of chronological information and radiocarbon calibration: the
566 program OxCal. *Archaeological Computing Newsletter*, 41, 11–16.

567 Bronk Ramsey, C. (1995). Radiocarbon calibration and analysis of stratigraphy: The OxCal program.
568 *Radiocarbon*, 37, 425–430. <https://doi.org/10.1017/S0033822200030903>

569 Bronk Ramsey, C. (2001). Development of the radiocarbon calibration program OxCal. *Radiocarbon*,
570 43, 355–363. <https://doi.org/10.1017/S0033822200038212>

571 Bronk Ramsey, C. (2008). Deposition models for chronological records. *Quaternary Science Reviews*,
572 27, 42–60. <https://doi.org/10.1016/j.quascirev.2007.01.019>

573 Bronk Ramsey, C., & Lee, S. (2013). Recent and planned developments of the program OxCal.
574 *Radiocarbon*, 55, 720–730. https://doi.org/10.2458/azu_js_rc.55.16215

575 Bronk Ramsey, C., van der Plicht, J., & Weninger, B. (2001). 'Wiggle matching' radiocarbon dates.
576 *Radiocarbon*, 43, 381–389. <https://doi.org/10.1017/S0033822200038248>

577 Bugmann, H., Brang, P., Elkin, C., Henne, P. D., Jakoby, O., Lévesque, M., ... Zimmermann, N. E. (2015).
578 Climate change impacts on tree species, forest properties, and ecosystem services. In C. C. Raible
579 & K. M. Strassmann (Eds.), *CH2014-impacts, toward quantitative scenarios of climate change*
580 *impacts in Switzerland* (pp. 79–88). Bern, Switzerland: OCCR, FOEN, MeteoSwiss, C2SM,
581 Agroscope, ProClim.

582 Colombaroli, D., Beckmann, M., van der Knaap, W. O., Curdy, P., & Tinner, W. (2013). Changes in
583 biodiversity and vegetation composition in the central Swiss Alps during the transition from
584 pristine forest to first farming. *Diversity and Distributions*, 19, 157–170.
585 <https://doi.org/10.1111/j.1472-4642.2012.00930.x>

586 Colombaroli, D., & Tinner, W. (2013). Determining the long-term changes in biodiversity and
587 provisioning services along a transect from Central Europe to the Mediterranean. *The Holocene*,
588 23, 1625–1634. <https://doi.org/10.1177/0959683613496290>

589 Conedera, M., Colombaroli, D., Tinner, W., Krebs, P., & Whitlock, C. (2017). Insights about past forest
590 dynamics as a tool for present and future forest management in Switzerland. *Forest Ecology and*
591 *Management*, 388, 100–112. <https://doi.org/10.1016/j.foreco.2016.10.027>

592 Conedera, M., Tinner, W., Cramer, S., Torriani, D., & Herold, A. (2006). Taxon-related pollen source
593 areas for lake basins in the southern Alps: an empirical approach. *Vegetation History and*
594 *Archaeobotany*, 15, 263–272. <https://doi.org/10.1007/s00334-006-0056-8>

595 Ellenberg, H. (2009). *Vegetation Ecology of Central Europe*. Cambridge, UK: University Press.

596 Feurdean, A., & Willis, K. J. (2008). Long-term variability of *Abies alba* in NW Romania: implications for
597 its conservation management. *Diversity and Distributions*, 14, 1004–1017.
598 <https://doi.org/10.1111/j.1472-4642.2008.00514.x>

- 599 Finsinger, W., & Tinner, W. (2005). Minimum count sums for charcoal-concentration estimates in
600 pollen slides: accuracy and potential errors. *The Holocene*, 15, 293–297.
601 <https://doi.org/10.1191/0959683605hl808rr>
- 602 Finsinger, W., & Tinner, W. (2006). Holocene vegetation and land-use changes in response to climatic
603 changes in the forelands of the southwestern Alps, Italy. *Journal of Quaternary Science*, 21, 243–
604 258. <https://doi.org/10.1002/jqs.971>
- 605 Fischer, J., Hafner, A., Stapfer, R., Marti, A., & Affolter, J. S. (2017). Neolithische Siedlungen in Nidau
606 am Bielersee. In *Archäologie Bern/Archéologie bernoise 2017* (pp. 126–155). Bern, Switzerland:
607 Jahrbuch des Archäologischen Dienstes des Kantons Bern.
- 608 FOREST EUROPE (2015). State of Europe's forests 2015.
- 609 Foster, D. R., Schoonmaker, P. K., & Pickett, S. T. A. (1990). Insights from paleoecology to community
610 ecology. *Trends in Ecology and Evolution*, 5, 119–122. [https://doi.org/10.1016/0169-](https://doi.org/10.1016/0169-5347(90)90166-B)
611 [5347\(90\)90166-B](https://doi.org/10.1016/0169-5347(90)90166-B)
- 612 van Geel, B., Mur, L. R., Ralska-Jasiewiczowa, M., & Goslar, T. (1994). Fossil akinetes of *Aphanizomenon*
613 and *Anabaena* as indicators of medieval phosphate-eutrophication of Lake Gosciarz (Central
614 Poland). *Review of Palaeobotany and Palynology*, 83, 97–105. [https://doi.org/10.1016/0034-](https://doi.org/10.1016/0034-6667(94)90061-2)
615 [6667\(94\)90061-2](https://doi.org/10.1016/0034-6667(94)90061-2)
- 616 Giesecke, T., Ammann, B., & Brande, A. (2014). Palynological richness and evenness: insights from the
617 taxa accumulation curve. *Vegetation History and Archaeobotany*, 23, 217–228.
618 <https://doi.org/10.1007/s00334-014-0435-5>
- 619 Gobet, E., & Tinner, W. (2012). Von der Ur- zur Kulturlandschaft. In A. Meyerhans (Ed.), *Geschichte des*
620 *Kantons Schwyz, Band 1: Zeiten und Räume, Frühzeit bis 1350* (pp. 37–57). Zurich, Switzerland:
621 Chronos.
- 622 Gobet, E., Tinner, W., Hochuli, P. A., van Leeuwen, J. F. N., & Ammann, B. (2003). Middle to Late
623 Holocene vegetation history in the Upper Engadine (Swiss Alps): the role of man and fire.
624 *Vegetation History and Archaeobotany*, 12, 143–163. [https://doi.org/10.1007/s00334-003-0017-](https://doi.org/10.1007/s00334-003-0017-4)
625 [4](https://doi.org/10.1007/s00334-003-0017-4)
- 626 Gobet, E., Vescovi, E., & Tinner, W. (2010). Ein paläoökologischer Beitrag zum besseren Verständnis
627 der natürlichen Vegetation der Schweiz. *Botanica Helvetica*, 120, 105–115.
628 <https://doi.org/10.1007/s00035-010-0080-2>
- 629 Green, D. G. (1981). Time-series and post-glacial forest ecology. *Quaternary Research*, 15, 265–277.
630 [https://doi.org/10.1016/0033-5894\(81\)90030-2](https://doi.org/10.1016/0033-5894(81)90030-2)
- 631 Gross-Klee, E., & Maise, C. (1997). Sonne, Vulkane und Seeufersiedlungen. *Jahrbuch der*
632 *Schweizerischen Gesellschaft für Ur- und Frühgeschichte*, 80, 85–94.
- 633 Guilaine, J. (2015). The neolithization of Mediterranean Europe: mobility and interactions from Near
634 East to the Iberian Peninsula. In C. Fowler, G. Bailey & D. Hofmann (Eds.), *The Oxford handbook*
635 *of Neolithic Europe* (pp. 81–98). Oxford, UK: Oxford University Press.
- 636 Guthruf, J., Zeh, M., & Guthruf-Seiler, K. (1999). *Kleinseen im Kanton Bern*. Bern, Switzerland: Haupt.
- 637 Haas, J. N., Karg, S., & Rasmussen, P. (1998). Beech leaves and twigs used as winter fodder: examples
638 from historic and prehistoric times. In M. Charles, P. Halstead & G. Jones (Eds.), *Fodder:*

639 *archaeological, historical and ethnographic studies* (pp. 81–86). Oxford, UK: Oxbow Books and
640 the Association for Environmental Archaeology.

641 Haas, J. N., Richoz, I., Tinner, W., & Wick, L. (1998). Synchronous Holocene climatic oscillations
642 recorded on the Swiss Plateau and at timberline in the Alps. *The Holocene*, 8, 301–309.
643 <https://doi.org/10.1191/095968398675491173>

644 Haas, J. N., & Schweingruber, F. (1993). Wood-anatomical evidence of pollarding in ash stems from the
645 Valais, Switzerland. *Dendrochronologia*, 11, 35–43.

646 Hadorn, P. (1992) *Vegetationsgeschichtliche Studie am Nordufer des Lac de Neuchâtel:*
647 *Pollenanalytische Untersuchungen im Loclat, in der Bucht von Hauterive/Saint-Blaise und in den*
648 *neolithischen Ufersiedlungen von Saint-Blaise/Bain des Dames*. PhD dissertation, University of
649 Bern, Bern, Switzerland.

650 Hafner, A., Harb, C., Amstutz, M., Francuz, J., & Moll-Dau, F. (2012). Moosseedorf, Moossee Oststation,
651 Strandbad. Strandbadneubau, Pfahlbauten und das älteste Boot der Schweiz. *Jahrbuch des*
652 *Archäologischen Dienstes des Kantons Bern*, 71–77.

653 Hafner, A., & Suter, P. J. (2000). -3400. *Die Entwicklung der Bauerngesellschaften im 4. Jahrtausend v.*
654 *Chr. am Bielersee aufgrund der Rettungsgrabungen von Nidau und Sutz-Lattrigen. Ufersiedlungen*
655 *am Bielersee (6th vol.)*. Bern, Switzerland: Berner Lehrmittel- und Medienverlag.

656 Harb, C. (2017). *Moosseedorf, Moossee. Ein Überblick über 160 Jahre Pfahlbauforschung*. Bern,
657 Switzerland: Archäologischer Dienst des Kantons Bern.

658 Heiri, O., Tinner, W., & Lotter, A. F. (2004). Evidence for cooler European summers during periods of
659 changing meltwater flux to the North Atlantic. *Proceedings of the National Academy of Sciences*
660 *of the United States of America*, 101, 15285–15288. <https://doi.org/10.1073/pnas.0406594101>

661 Henne, P. D., Elkin, C., Franke, J., Colombaroli, D., Calò, C., La Mantia, ... Tinner, W. (2015). Reviving
662 extinct Mediterranean forests increases ecosystem potential in a warmer future. *Frontiers in*
663 *Ecology and the Environment*, 13, 356–362. <https://doi.org/10.1890/150027>

664 Henne, P. D., Bigalke, M., Büntgen, U., Colombaroli, D., Conedera, M., Feller, U., ... Tinner, W. (2018).
665 An empirical perspective for understanding climate change impacts in Switzerland. *Regional*
666 *Environmental Change*, 18, 205–221. <https://doi.org/10.1007/s10113-017-1182-9>

667 Hillbrand, M., van Geel, B., Hasenfratz, A., Hadorn, P., & Haas, J. N. (2014). Non-pollen palynomorphs
668 show human- and livestock-induced eutrophication of Lake Nussbaumersee (Thurgau,
669 Switzerland) since Neolithic times (3840 BC). *The Holocene*, 24, 559–568.
670 <https://doi.org/10.1177/0959683614522307>

671 Hurlbert, S. H. (1971). The nonconcept of species diversity: a critique and alternative parameters.
672 *Ecology*, 52, 577–586. <https://doi.org/10.2307/1934145>

673 Jacomet, S., Ebersbach, R., Akeret, Ö., Antolín, F., Baum, T., Bogaard, A., ... Wick, L. (2016). On-site data
674 cast doubts on the hypothesis of shifting cultivation in the late Neolithic (c. 4300–2400 cal. BC):
675 Landscape management as an alternative paradigm. *The Holocene*, 26, 1858–1874.
676 <https://doi.org/10.1177/0959683616645941>

677 Juggins, S. (1991). Zone 1.2. freeware. DOS program for the zonation (constrained clustering) of
678 palaeoecological data. <https://www.staff.ncl.ac.uk/stephen.juggins/software.htm>.

679 Kalis, A. J., Merkt, J., & Wunderlich, J. (2003). Environmental changes during the Holocene climatic
680 optimum in central Europe – human impact and natural causes. *Quaternary Science Reviews*, 22,
681 33–79. [https://doi.org/10.1016/S0277-3791\(02\)00181-6](https://doi.org/10.1016/S0277-3791(02)00181-6)

682 Kaufmann, S., Hauck, M., & Leuschner, C. (2018). Effects of natural forest dynamics on vascular plant,
683 bryophyte, and lichen diversity in primeval *Fagus sylvatica* forests and comparison with
684 production forests. *Journal of Ecology*, <https://doi.org/10.1111/1365-2745.12981>

685 Kleinmann, A., Merkt, J., & Müller, H. (2015). Sedimente des Degersees: Ein Umweltarchiv –
686 Sedimentologie und Palynologie. In M. Mainberger, J. Merkt, & A. Kleinmann (Eds.),
687 *Pfahlbausiedlungen am Degersee. Archäologische und naturwissenschaftliche Untersuchungen*
688 (pp. 409–471). Darmstadt, Germany: Theiss.

689 van der Knaap, W. O. (2009). Estimating pollen diversity from pollen accumulation rates: A method to
690 assess taxonomic richness in the landscape. *The Holocene*, 19, 159–163.
691 <https://doi.org/10.1177/0959683608098962>

692 van der Knaap, W. O., van Leeuwen, J. F. N., & Ammann, B. (2004). The first rise and fall of *Fagus*
693 *sylvatica* and interactions with *Abies alba* at Faulenseemoos (Swiss Plateau) 6,900–6,000 cal yr
694 BP. *Acta Palaeobotanica*, 44, 249–266.

695 Kovats, R. S., Valentini, R., Bouwer, L. M., Georgopoulou, E., Jacob, D., Martin, E., ... Soussana, J.-F.
696 (2014). Europe. In V. R. Barros, C. B. Field, D. J. Dokken, M. D. Mastrandrea, K. J. Mach, T. E. Bilir
697 ... L. L. White (Eds.), *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B:*
698 *Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*
699 *Intergovernmental Panel on Climate Change* (pp. 1267–1326). Cambridge, UK and New York, NY:
700 Cambridge University Press.

701 Lindbladh, M., Niklasson, M., Karlsson, M., Björkman, L., & Churski, M. (2008). Close anthropogenic
702 control of *Fagus sylvatica* establishment and expansion in a Swedish protected landscape –
703 implications for forest history and conservation. *Journal of Biogeography*, 35, 682–697.
704 <https://doi.org/10.1111/j.1365-2699.2007.01813.x>

705 Lotter, A. F. (1999). Late-glacial and Holocene vegetation history and dynamics as shown by pollen and
706 plant macrofossil analyses in annually laminated sediments from Soppensee, central Switzerland.
707 *Vegetation History and Archaeobotany*, 8, 165–184. <https://doi.org/10.1007/BF02342718>

708 Magny, M. (2004). Holocene climate variability as reflected by mid-European lake-level fluctuations
709 and its probable impact on prehistoric human settlements. *Quaternary International*, 113, 65–
710 79. [https://doi.org/10.1016/S1040-6182\(03\)00080-6](https://doi.org/10.1016/S1040-6182(03)00080-6)

711 Magny, M. (2013). Orbital, ice-sheet, and possible solar forcing of Holocene lake-level fluctuations in
712 west-central Europe: a comment on Bleicher. *The Holocene*, 23, 1202–1212.
713 <https://doi.org/10.1177/0959683613483627>

714 McKinney, L. V., Nielsen, N. R., Hansen J. K., & Kjær E. D. (2011). Presence of natural genetic resistance
715 in *Fraxinus excelsior* (Oleaceae) to *Chalara fraxinea* (Ascomycota): an emerging infectious
716 disease. *Heredity*, 106, 788–797. <https://doi.org/10.1038/hdy.2010.119>

717 MeteoSwiss (2017). Klimadiagramme und Normwerte pro Station. Zurich-Airport, Switzerland: Federal
718 Office of Meteorology and Climatology MeteoSwiss.

719 Moore, P. D., Webb, J. A., & Collison, M. E. (1991). *Pollen Analysis*. Oxford, UK: Blackwell Scientific
720 Publications.

- 721 Müller, J. (2015). Movement of plants, animals, ideas, and people in South-East Europe. In C. Fowler,
722 G. Bailey, & D. Hofmann (Eds.), *The Oxford handbook of Neolithic Europe* (pp. 63–80). Oxford,
723 UK: Oxford University Press.
- 724 Odgaard, B. V. (1999). Fossil pollen as a record of past biodiversity. *Journal of Biogeography*, 26, 7–17.
725 <https://doi.org/10.1046/j.1365-2699.1999.00280.x>
- 726 Pott, R. (1985). Vegetationsgeschichtliche und pflanzensoziologische Untersuchungen zur
727 Niederwaldwirtschaft in Westfalen. *Abhandlungen aus dem Westfälischen Museum für*
728 *Naturkunde*, 47, 1–78.
- 729 Puhe, J., & Ulrich, B. (2001). *Global climate change and human impacts on forest ecosystems –*
730 *postglacial development, present situation, and future trends in Central Europe*. Berlin, Germany:
731 Springer.
- 732 R Development Core Team (2016). *R: A language and environment for statistical computing*. R
733 Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- 734 Reille, M. (1992). *Pollen et spores d'Europe et d'Afrique du Nord*. Marseille, France: Laboratoire de
735 botanique historique et palynologie.
- 736 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., ... van der Plicht J
737 (2013). IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP.
738 *Radiocarbon*, 55, 1869–1887. https://doi.org/10.2458/azu_js_rc.55.16947
- 739 Renner, E. (1982). *Beiträge zur Gletschergeschichte des Gotthardgebietes und dendroklimatologische*
740 *Analysen an fossilen Hölzern*. Physische Geographie, Vol. 8. University of Zurich, Switzerland.
- 741 Rey, F., Gobet, E., van Leeuwen, J. F. N., Gilli, A., van Raden, U. J., Hafner, A., ... Tinner, W. (2017).
742 Vegetational and agricultural dynamics at Burgäschisee (Swiss Plateau) recorded for 18,700 years
743 by multi-proxy evidence from partly varved sediments. *Vegetation history and archaeobotany*,
744 26, 571–586. <https://doi.org/10.1007/s00334-017-0635-x>
- 745 Rey, F., Gobet, E., Szidat, S., Lotter, A. F., Gilli, A., Hafner, A., & Tinner, W. (2018). Radiocarbon wiggle
746 matching on laminated sediments delivers high-precision chronologies. *Radiocarbon*,
747 <https://doi.org/10.1017/RDC.2018.47>
- 748 Rey, F., Schwörer, C., Gobet, E., Colombaroli, D., van Leeuwen, J. F. N., Schleiss, S., & Tinner, W. (2013).
749 Climatic and human impacts on mountain vegetation at Lauenensee (Bernese Alps, Switzerland)
750 during the last 14,000 years. *The Holocene*, 23, 1415–1427.
751 <https://doi.org/10.1177/0959683613489585>
- 752 Rösch, M., Kleinmann, A., Lechterbeck, J., & Wick, L. (2014a). Botanical offsite and on-site data as
753 indicators of different land use systems: a discussion with examples from Southwest Germany.
754 *Vegetation History and Archaeobotany*, 23, 121–133. [https://doi.org/10.1007/s00334-014-0437-](https://doi.org/10.1007/s00334-014-0437-3)
755 3
- 756 Rösch, M., Kleinmann, A., Lechterbeck, J., & Wick, L. (2014b). Erratum to: Botanical off-site and on-site
757 data as indicators of different land use systems: a discussion with examples from Southwest
758 Germany. *Vegetation History and Archaeobotany*, 23, 647–648. [https://doi.org/10.1007/s00334-](https://doi.org/10.1007/s00334-014-0477-8)
759 014-0477-8
- 760 Rösch, M., & Lechterbeck, J. (2016). Seven Millennia of human impact as reflected in high resolution
761 pollen profile from the profundal sediments of Litzelsee, Lake Constance region, Germany.

762 *Vegetation History and Archaeobotany*, 25, 339–358. [https://doi.org/10.1007/s00334-015-0552-](https://doi.org/10.1007/s00334-015-0552-9)
763 9

764 Ruosch, M., Spahni, R., Joos, F., Henne, P. D., van der Knaap, W.O., & Tinner, W. (2016). Past and future
765 evolution of *Abies alba* forests in Europe – comparison of a dynamic vegetation model with
766 palaeo data and observations. *Global Change Biology*, 22, 727–740.
767 <https://doi.org/10.1111/gcb.13075>

768 Schier, W. (2015). Central and Eastern Europe. In C. Fowler, G. Bailey & D. Hofmann (Eds.), *The Oxford*
769 *handbook of Neolithic Europe* (pp. 99–120). Oxford, UK: Oxford University Press.

770 Schwörer, C., Colombaroli, D., Kaltenrieder, P., Rey, F., & Tinner, W. (2015). Early human impact
771 (5,000–3,000 BC) affects mountain forest dynamics in the Alps. *Journal of Ecology*, 103, 281–295.
772 <https://doi.org/10.1111/1365-2745.12354>

773 Steiger, P. (2010). *Wälder der Schweiz*. Bern, Switzerland: Ott.

774 Steinhilber, F., Beer, J., & Fröhlich, C. (2009). Total solar irradiance during the Holocene. *Geophysical*
775 *Research Letters*, 36, L19704. <https://doi.org/10.1029/2009GL040142>

776 Stöckli, W. E. (2016). *Urgeschichte der Schweiz im Überblick (15,000 v. Chr. – Christi Geburt). Die*
777 *Konstruktion einer Urgeschichte*. Basel, Switzerland: Veröffentlichung der Archäologie Schweiz.

778 Stockmarr, J. (1971). Tablets with spores used in absolute pollen analysis. *Pollen et spores*, 13, 615–
779 621.

780 Sugita, S. (1994). Pollen representation of vegetation in Quaternary sediments: theory and method in
781 patchy vegetation. *Journal of Ecology*, 82, 881–897.

782 Suter, P. J. (2017). *Um 2700 v. Chr. Wandel und Kontinuität in den Ufersiedlungen am Bielersee*. Bern,
783 Switzerland: Archäologischer Dienst des Kantons Bern.

784 Systat (2007). *Mystat 12*. San Jose, CA: Systat Software Inc.

785 Tinner, W., & Ammann, B. (2005). Long-term responses of mountain ecosystems to environmental
786 changes: Resilience, adjustment and vulnerability. In U. M. Huber, H. Bugmann & M. Reasoner
787 (Eds.), *Global change and mountain research – state of knowledge overview* (pp. 133–144).
788 Dordrecht, The Netherlands: Springer.

789 Tinner, W., Beer, R., Bigler, C., Clegg, B. F., Jones, R. T., Kaltenrieder, P., ... Hu, F. S. (2015). Late-
790 Holocene climate variability and ecosystem responses in Alaska inferred from high-resolution
791 multiproxy sediment analyses at Grizzly Lake. *Quaternary Science Reviews*, 126, 41–56.
792 <https://doi.org/10.1016/j.quascirev.2015.08.019>

793 Tinner, W., Colombaroli, D., Heiri, O., Henne, P. D., Steinacher, M., Untenecker, J., ... Valsecchi, V.
794 (2013). The past ecology of *Abies alba* provides new perspectives on future responses of silver
795 fir forests to global warming. *Ecological Monographs*, 83, 419–439. [https://doi.org/10.1890/12-](https://doi.org/10.1890/12-2231.1)
796 2231.1

797 Tinner, W., Conedera, M., Ammann, B., & Lotter, A. F. (2005). Fire ecology north and south of the Alps
798 since the last ice age. *The Holocene*, 15, 1214–1226.
799 <https://doi.org/10.1191/0959683605hl892rp>

800 Tinner, W., Conedera, M., Ammann, B., Gaggeler, H. W., Gedy, S., Jones, R. & Sagesser, B. (1998).
801 Pollen and charcoal in lake sediments compared with historically documented forest fires in

802 southern Switzerland since AD 1920. *The Holocene*, 8, 31–42.
803 <https://doi.org/10.1191/095968398667205430>

804 Tinner, W., Conedera, M., Gobet, E., Hubschmid, P., Wehrli, M., & Ammann, B. (2000). A
805 palaeoecological attempt to classify fire sensitivity of trees in the southern Alps. *The Holocene*,
806 10, 565–574. <https://doi.org/10.1191/095968300674242447>

807 Tinner, W., & Hu, F. S. (2003). Size parameters, size-class distribution and area number relationship of
808 microscopic charcoal: relevance for fire reconstruction. *The Holocene*, 13, 499–505.
809 <https://doi.org/10.1191/0959683603hl615rp>

810 Tinner, W., Hubschmid, P., Wehrli, M., Ammann, B., & Conedera, M. (1999). Long-term forest fire
811 ecology and dynamics in southern Switzerland. *Journal of Ecology*, 87, 273–289.
812 <https://doi.org/10.1046/j.1365-2745.1999.00346.x>

813 Tinner, W., van Leeuwen, J. F. N., Colombaroli, D., Vescovi, E., van der Knaap, W. O., Henne, P. D., ... La
814 Mantia, T. (2009). Holocene environmental changes at Gorgo Basso, a coastal lake in southern
815 Sicily, Italy. *Quaternary Science Reviews*, 28, 1498–1510.
816 <https://doi.org/10.1016/j.quascirev.2009.02.001>

817 Tinner, W., & Lotter, A. F. (2006). Holocene expansions of *Fagus silvatica* and *Abies alba* in Central
818 Europe: where are we after eight decades of debate? *Quaternary Science Reviews*, 25, 526–549
819 <https://doi.org/10.1016/j.quascirev.2005.03.017>

820 Tinner, W., Lotter, A. F., Ammann, B., Conedera, M., Hubschmid, P., van Leeuwen, J. F. N., & Wehrli,
821 M. (2003). Climatic change and contemporaneous land-use phases north and south of the Alps
822 2,300 BC to 800 AD. *Quaternary Science Reviews*, 22, 1447–1460.
823 [https://doi.org/10.1016/S0277-3791\(03\)00083-0](https://doi.org/10.1016/S0277-3791(03)00083-0)

824 Valsecchi, V., Carraro, G., Conedera, M., & Tinner, W. (2010). Late-Holocene vegetation and land-use
825 dynamics in the Southern Alps (Switzerland) as a basis for nature protection and forest
826 management. *The Holocene*, 20, 483–495. <https://doi.org/10.1177/0959683609355178>

827 Vinther, B. M., Clausen, H. B., Johnson, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., ...
828 Heinemeier, J. (2006). A synchronized dating of three Greenland ice cores throughout the
829 Holocene. *Journal of Geophysical Research*, 111, D13102.
830 <https://doi.org/10.1029/2005JD006921>

831 Wehrli, M., Tinner, W., & Ammann, B. (2007). 16,000 years of vegetation and settlement history from
832 Egelsee (Menzingen, central Switzerland). *The Holocene*, 17, 747–761.
833 <https://doi.org/10.1177/0959683607080515>

834 Wey, O. (2012). *Die Cortaillod-Kultur am Burgäschisee: Materialvorlage und Synthese zu den*
835 *neolithischen Fundkomplexen von Burgäschisee-Ost, -Südwest, -Süd und -Nord*. Bern,
836 Switzerland: Stämpfli.

837 Willis, K. J., & Birks, H. J. B. (2006). What is natural? The need for a long-term perspective in biodiversity
838 conservation. *Science*, 314, 1261–1265. <https://doi.org/10.1126/science.1122667>




FIGURE 1 Overview map of Switzerland and the neighbouring countries with the two sites of this study (in red) and other discussed sites. 1 Moossee; 2 Burgäschisee; 3 Lobsigensee (Ammann, 1989); 4 Lake Biel (Fischer, Hafner, Stapfer, Marti, & Affolter, 2017; Hafner & Suter, 2000; Stöckli, 2016; Suter 2017); 5 Soppensee (Lotter, 1999); 6 Zürich Opéra (Bleicher et al., 2017); 7 Egelsee (Wehrli, Tinner, & Ammann, 2007); 8 Mindelsee (Rösch, Kleinmann, Lechterbeck, & Wick, 2014a, b); 9 Degersee (Kleinmann, Merkt, & Müller, 2015); 10 Lago di Origlio (Tinner, Hubschmid, Wehrli, Ammann, & Conedera, 1999). A = Austria, D = Germany, F = France, FL = Liechtenstein, I = Italy (topographic map: © swisstopo)




FIGURE 2 Age-depth models of (a) Moossee and (b) Burgäschisee. Black dots show the calibrated ages with 95% (2σ) probabilities (IntCal13, Reimer et al., 2013). The black lines are the modelled chronologies (OxCal; Bronk Ramsey, 1994, 1995, 2001, 2008; Bronk Ramsey, van der Plicht, & Weninger, 2001; Bronk Ramsey & Lee, 2013). One age at Burgäschisee (dark grey x) was treated as an outlier. The grey lines indicate the 95% (2σ) probabilities of the models.




FIGURE 3 Moossee high-resolution sequence. Presented are selected pollen, spore, algae and cyanobacteria percentages, microscopic charcoal influx values, macroscopic charcoal concentrations (concentration = number of particles $\geq 600 \mu\text{m}/17.5 \text{ cm}^3$, black bars, plotted on top axis), selected plant macrofossil concentrations (concentration = number of remains/ 17.5 cm^3 , dark green bars, plotted on top axis), presence of *Abies alba* stomata (dark green dots), palynological richness (PRI, light blue), evenness-detrended palynological richness (DE-PRI, red) as well as palynological evenness (PIE). Empty curves show 10 \times exaggerations. Light grey bars indicate the time windows of local archaeological findings reflecting settlement phases (1–6). BS = bud scales, C = cones, F = fruits, N = needles, S = seeds.




FIGURE 4 Burgäschisee high-resolution sequence. Presented are selected pollen, spore, algae and cyanobacteria percentages, microscopic charcoal influx values, macroscopic charcoal concentrations (concentration = number of particles $\geq 600 \mu\text{m}^3$ (standard volume), black bars, plotted on top axis), selected plant macrofossil concentrations (concentration = number of remains/11 cm^3 (standard volume), dark green bars, plotted on top axis), presence of *Abies alba* stomata (dark green dots), palynological richness (PRI, light blue), evenness-detrended palynological richness (DE-PRI, red) as well as palynological evenness (PIE). Empty curves show 10 \times exaggerations. Light grey bars indicate the time windows of local archaeological findings reflecting settlement phases (1–4). B = bud, BS = bud scales, CS = cone scales, F = fruits, FS = fruit scales, L = leaves, N = needles, S = seeds.




FIGURE 5 Cross-correlation analysis of microscopic charcoal influx vs. percentages of selected pollen types, palynological richness (PRI), palynological evenness (PIE) and evenness-detrended palynological richness (DE-PRI) from (a) Moossee and (b) Burgäschisee (6,500–4,150 cal year BP). 1 lag = 11.1 ± 0.2 cal years (Moossee), 8.7 ± 0.6 cal years (Burgäschisee). The solid black lines mark the significance level ($P < 0.05$).




FIGURE 6 Correlograms showing the correlation coefficients at lag 0 between microscopic charcoal influx and selected pollen types, spore types, palynological richness (PRI), palynological evenness (PIE) and evenness-detrended palynological richness (DE-PRI) for (a) Moossee and (b) Burgäschisee (6,500–4,150 cal year BP). The dotted lines mark the significance level ($P < 0.05$).




FIGURE 7 Cross-correlation analysis of percentages of selected pollen types, palynological richness (PRI) and evenness-detrended palynological richness (DE-PRI) from Moossee (Moos) vs. percentages of selected pollen types, PRI and DE-PRI from Burgäschisee (Burg) (6,500–4,150 cal year BP). 1 lag = 11.1 ± 0.2 cal. years. The solid black lines mark the significance level ($P < 0.05$).




FIGURE 8 Tree pollen percentages from selected sites (see Figure 1). Dark green bars indicate periods with fairly closed forests (tree pollen % >80). Light green bars indicate phases with forest openings (tree pollen % <80).




FIGURE 9 Percentages of selected pollen types, palynological richness (PRI), microscopic charcoal influx, macroscopic charcoal presence (≥600 µm) and local archaeological findings at Moossee and Burgäschisee compared to palaeoclimatic proxies. Purple graphs = Moossee, green graphs = Burgäschisee. Cultural indicators = \sum Cerealia-type + *Plantago lanceolata* + *Linum usitatissimum*-type. Bottom labelling, center: (a) Local archaeological settlement phases within the city of Zurich (Bleicher et al., 2017). (b) Dendrochronologically dated settlement phases at Lake Biel (Fischer, Hafner, Stapfer, Marti, & Affolter, 2017; Hafner & Suter, 2000; Stöckli, 2016; Suter 2017). Dark coloured bars (archaeology) indicate dendrochronologically dated settlement phases (Moossee, Burgäschisee and Zurich). (c) δTSI (total solar irradiance) (Steinhilber, Beer, & Fröhlich, 2009). (d) δ¹⁸O NGRIP (Vinther et al., 2006) with smoothing (loess = 0.05, red line). (e) Calibrated tree ring data from the Swiss Alps (Bircher, 1982, 1986; Renner, 1982). Dotted black lines mark data gaps (f) Periods of high lake levels in Eastern France (F) and Western Switzerland (CH) (Magny, 2004, 2013).